


Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

A292.9
S03W
Log 2

**FEDERAL-STATE-PRIVATE
COOPERATIVE SNOW SURVEYS**



U. S. DEPT. OF AGRICULTURE
NATIONAL AGRICULTURAL LIBRARY
RECEIVED

MAY 31 1972

PROCUREMENT SECTION
CURRENT SERIAL RECORDS

***WATER SUPPLY OUTLOOK
FOR
WESTERN UNITED STATES***

Including Columbia River Drainage in Canada

Prepared by

U. S. DEPARTMENT of AGRICULTURE ★ SOIL CONSERVATION SERVICE

Collaborating with
CALIFORNIA DEPARTMENT of WATER RESOURCES
and

BRITISH COLUMBIA DEPARTMENT of
LANDS, FORESTS and WATER RESOURCES

AS OF
MAY 1, 1972

TO RECIPIENTS OF WATER SUPPLY OUTLOOK REPORTS:

Most of the usable water in western states originates as mountain snowfall. This snowfall accumulates during the winter and spring, several months before the snow melts and appears as streamflow. Since the runoff from precipitation as snow is delayed, estimates of snowmelt runoff can be made well in advance of its occurrence. Streamflow forecasts published in this report are based principally on measurement of the water equivalent of the mountain snowpack.

Forecasts become more accurate as more of the data affecting runoff are measured. All forecasts assume that climatic factors during the remainder of the snow accumulation and melt season will interact with a resultant average effect on runoff. Early season forecasts are therefore subject to a greater change than those made on later dates.

The snow course measurement is obtained by sampling snow depth and water equivalent at surveyed and marked locations in mountain areas. A total of about ten samples are taken at each location. The average of these are reported as snow depth and water equivalent. These measurements are repeated in the same location near the same dates each year.

Snow surveys are made monthly or semi-monthly from January 1 through June 1 in most states. There are about 1900 snow courses in Western United States and in the Columbia Basin in British Columbia. Networks of automatic snow water equivalent and related data sensing devices, along with radio telemetry are expanding and will provide a continuous record of snow water and other parameters of key locations.

Detailed data on snow course and soil moisture measurements are presented in state and local reports. Other data on reservoir storage, summaries of precipitation, current streamflow, and soil moisture conditions at valley elevations are also included. The report for Western United States presents a broad picture of water supply outlook conditions, including selected streamflow forecasts, summary of snow accumulation to date, and storage in larger reservoirs.

Snow survey and soil moisture data for the period of record are published by the Soil Conservation Service by states about every five years. Data for the current year is summarized in a West-wide basic data summary and published about October 1 of each year.

COVER PHOTO NUMBER ORC 221-3

PUBLISHED BY SOIL CONSERVATION SERVICE

The Soil Conservation Service publishes reports following the principal snow survey dates from January 1 through June 1 in cooperation with state water administrators, agricultural experiment stations and others. Copies of the reports for Western United States and all state reports may be obtained from Soil Conservation Service, Western Regional Technical Service Center, Room 209, 701 N. W. Glisan, Portland, Oregon 97209.

Copies of state and local reports may also be obtained from state offices of the Soil Conservation Service in the following states:

STATE	ADDRESS
Alaska	P. O. Box "F", Palmer, Alaska 99645
Arizona	6029 Federal Building, Phoenix, Arizona 85025
Colorado (N. Mex.)	P. O. Box 17107, Denver, Colorado 80217
Idaho	Room 345, 304 N. 8th. St., Boise, Idaho 83702
Montana	P. O. Box 970, Bozeman, Montana 59715
Nevada	P. O. Box 4850, Reno Nevada 89505
Oregon	1218 S. W. Washington St., Portland, Oregon 97205
Utah	4012 Federal Bldg., 125 South State St., Salt Lake City, Utah 84111
Washington	360 U.S. Court House, Spokane, Washington 99201
Wyoming	P. O. Box 2440, Casper, Wyoming 82601

PUBLISHED BY OTHER AGENCIES

Water Supply Outlook reports prepared by other agencies include a report for California by the Water Supply Forecast and Snow Surveys Unit, California Department of Water Resources, P. O. Box 388, Sacramento, California 95802 --- and for British Columbia by the Department of Lands, Forests and Water Resources, Water Resources Service, Parliament Building, Victoria, British Columbia



WATER SUPPLY OUTLOOK FOR WESTERN UNITED STATES

Including Columbia River Drainage in Canada

ISSUED

MAY 1, 1972

The Soil Conservation Service coordinates snow surveys conducted by its staff and many cooperators, including the Bureau of Reclamation, Corps of Engineers, Forest Service, National Park Service, NOAA, National Weather Service, Geological Survey, and other Federal Agencies, Departments of State Government, Irrigation Districts, Power Companies, and others.

The Department of Water Resources coordinates snow surveys in California.


The Water Resources Service, Department of Lands, Forests, and Water Resources directs snow surveys in British Columbia.

This report was prepared by the Water Supply Forecasting Unit, Engineering Division, Soil Conservation Service, from data supplied by Snow Survey Supervisors of the Soil Conservation Service in the States of Alaska, Arizona, Colorado and New Mexico, Idaho, Montana, Nevada, Oregon, Utah, Washington, and Wyoming.

Data from California was supplied by the Chief, Water Supply Forecast and Snow Survey Unit, Department of Water Resources.

Data from British Columbia was supplied by the Chief, Hydrology Division, Water Investigations Branch, Department of Lands, Forests and Water Resources.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE
KENNETH E. GRANT, ADMINISTRATOR



PROSPECTIVE STREAMFLOW

WATER SUPPLY OUTLOOK

1972 SNOWMELT SEASON
MAY 1, 1972

HEAVY SNOWPACKS INDICATE STRONG PROBABILITY OF HIGH WATER PROBLEMS IN PARTS OF THE COLUMBIA, MISSOURI, UPPER GREEN RIVER, GREAT SALT LAKE DRAINAGES, AND IN ALASKA. SHORTAGES ANTICIPATED FOR WATER USERS ON NATURAL FLOW RIGHTS IN ARIZONA, NEW MEXICO, AND LIMITED AREAS IN SOUTHERN SECTIONS OF COLORADO, UTAH, NEVADA AND CALIFORNIA. RESERVOIR STORAGE GENERALLY ADEQUATE EXCEPT IN NEW MEXICO.

During April the southern areas of the West experienced a continuation of the dry weather which has prevailed since the beginning of 1972, intensifying the poor outlook for streamflow this summer. In northern areas temperatures were generally cool while precipitation was more typical, with areas of high and low precipitation.

The cool weather in northern areas delayed snowmelt and maintained a serious flood potential on many watersheds. If the weather during the May-June period consists of alternate warm and cool spells, or if it remains cool well into July as it did last year, high water problems from the major snowpack will be held to a minimum. However, if the weather turns warm for an extended period, and particularly if it is accompanied by warm rains, high water damage could become severe in areas above reservoirs, as well as below any not holding adequate flood control space.

The California Department of Water Resources reports that, although above normal precipitation during April through the central Sierra provided some improvement there, elsewhere the State's snowmelt runoff potential continued to deteriorate. The snowpack in California's Cascade Mountains and the Sierra Nevada Ranges on May 1 was about 70 percent of that normally expected for that date. While streamflow during April was below normal in all areas, the total storage in the State's major reservoirs on May 1 was near normal. Southern California, for the fourth consecutive month, experienced much below average precipitation.

Canadian tributaries to the Columbia River have maximum or near maximum of record snowpacks, according to the British Columbia Water Resources Service, Department of Lands, Forests and Water Resources. On these watersheds the snow varies from 141 percent for the Columbia below Revelstoke to 199 percent on the Similkameen River. Runoff from these streams is also expected to be maximum or near maximum of record.

In the Columbia Basin the snow is not only excessive in British Columbia, but is also very heavy on Montana's upper Clark Fork and Bitterroot rivers, on the Cascade Mountains in Washington, on Idaho's Clearwater and Bruneau rivers and on Oregon's Hood, Owyhee, Malheur and Umatilla rivers. Snow here ranges from near 165 to over 200 percent of average. Snow on the majority of watersheds in the Columbia Basin is over 140 percent of normal. Most streams will yield from near 125 to 165 percent of usual amounts.

An unregulated peak flow of near 850 thousand to one million second feet is expected at The Dalles, Oregon. These flows would result in a river stage at Vancouver of 28 to 31 feet. Actual flow and river stage will depend on snowmelt conditions during late May and early June, and on the amount that river regulation reduces the flow. River management agencies have indicated that reservoir regulation will reduce the river stage at Vancouver to about 21 to 23 feet.

In Alaska snowmelt runoff of the Chena and Salcha rivers is forecast at 170 and 146 percent of normal, respectively. These forecasts are typical of conditions in much of Alaska.

In the Missouri Basin snowpacks are greatest, with respect to normal, along the Continental Divide between Glacier and Yellowstone National Parks. Here some areas show maximum of record snow. In general, the Montana snowpack is 130 to 150 percent. In Wyoming it is near 120 to 150 percent except on the North Platte where it is a little above average. Most Montana streams will yield 30 to 60 percent above normal amounts.

In Wyoming, except for the North Platte and its tributaries and the Little Snake River near the Colorado border, where streamflow should be near average, forecasts for most other streams in the Missouri, Snake, Green and Bear river basins range from near 120 to 160 percent of usual amounts.

SUMMARY OF SNOW WATER EQUIVALENT MEASUREMENTS

MAY 1, 1972

MAJOR BASIN AND SUB - WATERSHED	WATER EQUIVALENT IN PERCENT OF :		MAJOR BASIN AND SUB - WATERSHED	WATER EQUIVALENT IN PERCENT OF :	
	LAST YEAR	AVERAGE		LAST YEAR	AVERAGE
MISSOURI BASIN			SNAKE BASIN		
Jefferson	102	154	SNAKE above Jackson, Wyo.	91	144
Madison	85	126	SNAKE above Hiese, Idaho	81	153
Gallatin	74	109	SNAKE abv. American Falls Res.	88	150
Missouri Main Stem	104	138	Henry's Fork	70	120
Yellowstone	82	127	Southern Idaho Tributaries	105	165
Shoshone	82	132	Big and Little Wood	85	125
Wind	76	147	Boise	105	160
North Platte	69	116	Owyhee	290	215
South Platte	72	97	Payette	90	150
			Malheur	105	190
ARKANSAS BASIN			Weiser	75	135
Arkansas	93	88	Burnt	85	145
Cucharas-Purgatoire	0	0	Powder	100	155
			Salmon	105	150
RIO GRANDE BASIN			Grande Ronde	115	150
Rio Grande (Colo.)	99	60	Clearwater	145	180
Rio Grande abv. Otowi Bridge	93	53			
Pecos	0	0	LOWER COLUMBIA BASIN		
			Yakima	98	214
COLORADO BASIN			Umatilla	141	177
Green (Wyo.)	88	152	John Day	100	155
Yampa - White	58	77	Deschutes - Crooked	95	145
Duchesne	97	92	Hood	95	180
Price	64	58	Willamette	80	145
Upper Colorado	76	93	Lewis	76	182
Gunnison	70	63	Cowlitz	106	173
San Juan	102	52			
Dolores	29	15	PACIFIC COASTAL BASIN		
Virgin	93	51	Puget Sound	91	145
Gila	0	0	Olympic Peninsula	84	130
Salt	0	0	Umpqua - Rogue	75	145
			Klamath	105	130
GREAT BASIN			Trinity	42	70
Bear	91	154			
Logan	94	169	CALIFORNIA		
Ogden	103	176	CENTRAL VALLEY		
Weber	110	128	Upper Sacramento	56	90
Provo - Utah Lake	82	68	Feather	41	70
Jordan	118	134	Yuba	51	95
Sevier	54	42	American	59	80
Walker - Carson	81	101	Mokelumne	68	75
Tahoe - Truckee	61	103	Stanislaus	67	70
Humboldt	85	120	Tuolumne	74	70
Lake Co. (Oregon)	75	115	Merced	78	70
Harney Basin (Oregon)	90	145	San Joaquin	67	60
			Kings	65	55
UPPER COLUMBIA BASIN			Kaweah	50	40
Columbia (Canada)	123	142	Tule	10	5
Kootenai	139	156	Kern	23	15
Clark Fork	130	160			
Bitterroot	127	167	Data for California Watersheds supplied by Dept. of Water Resources, and for British Columbia Watersheds by Dept. of Lands, Forests and Water Resources.		
Flathead	130	148			
Spokane	135	155	Average is for 1953-67 period. California averages are for the period 1931-70. Based on Selected Snow Courses determined by Dis- tribution within the Basin, Length of Record and Repetitive Monthly Measurement Schedules.		
Okanogan	145	187			
Methow	105	184			
Chelan	124	163			
Wenatchee	117	238			

Colorado's South Platte will produce a little below normal amounts, but has excellent reservoir storage. Outlook for the Arkansas is not so favorable. Here reservoir storage is poor and streamflow prospects are for near one-third less than normal.

Streamflow prospects are very unfavorable in Arizona and New Mexico where practically all of the snow has gone and most streams are already dropping to base flow values. Residual streamflow in Arizona will be near 15 to 35 percent average, while in New Mexico it will range from about 30 to 65 percent. Reservoir storage on Arizona's Salt River Project will be adequate, but New Mexico reservoir water is short. Storage in Conchas Reservoir is 45 percent, in Elephant Butte 53 percent, in El Vado 26 percent of the normal amounts.

In the Upper Colorado Basin the water outlook is excellent in Wyoming, fair to good in Utah's Uinta Basin and on Colorado's White, Yampa and upper Colorado rivers. It is poor for the Gunnison, Dolores, San Juan and smaller streams in southern Utah. Reservoir storage is favorable.

Inflow to Lake Powell on the Colorado River is forecast at 86 percent.

In the Great Basin reservoir storage is very favorable. This will be particularly important in Nevada and southern Utah. While flow of the Humboldt River will be near average, streams draining into Nevada from the Sierra Mountains will be near three-fourths of the normal amount. Late season shortages are expected in southern Utah and southern Nevada for water users dependent on natural streamflow.

With the principal exception of New Mexico, water stored for irrigation purposes continues near or well above average in the West. Storage space reserved for flood control operations has been greatly increased as reservoirs have been drawn down sharply to provide room for expected high runoff in many of the high snowpack areas.

MISSOURI BASIN

Snow along the Continental Divide from Glacier National Park to Yellowstone National Park continues heavy, with some areas having a maximum of record snowpack. The snow decreases to near average on the Gallatin River and in the Snowy and Crazy Mountains. On the Yellowstone and Upper Clark's Fork rivers, as well as on interior mountain ranges in Montana, the snow is well above average. In general, the Montana snowpack is 130 to 150 percent average.

Streams in the heavy snowpack areas of

Montana, particularly those with headwaters along the Continental Divide, are forecast to yield near 30 to 60 percent above average amounts. In the southwestern portion of the state, streams should yield a little less runoff than last year, while streams farther north are expected to produce more runoff. Other streams in the state will yield near average to 30 percent above average amounts.

In Wyoming the snow is well above average, but near 20 to 25 percent less than the record or near record amounts measured last year. It is near 120 to 150 percent of the normal snowpack. Stream forecasts range from near 120 to 145 percent of usual amounts for the Clark's Fork, Shoshone, Wind and Bighorn rivers, as well as for smaller streams such as the Tongue, Powder and other streams draining from the Big Horn Mountains.

On the North Platte River system, prospective streamflow is generally within 10 percent of average amounts, but ranges from 90 percent on the Laramie River near Glendevy to 127 percent on the Little Laramie near Fillmore.

Forecasts for the South Platte River and its tributaries are about the same as expected last month. They range from 86 percent to 98 percent.

Reservoir storage is excellent. This, combined with prospective streamflow provides a good to excellent water outlook for the coming summer. If the early summer months are dry, some limited shortages may develop for water users on natural flow rights who are served by streams draining from the Black Hills. However, reservoir storage here is good, as illustrated by Belle Fourche which holds 148 percent of its usual amount.

ARKANSAS BASIN

Due to continuing dry weather, streamflow forecasts have been steadily dropping since February. Forecasts for streams in Colorado now range from 60 percent of average for the Purgatoire at Trinidad to 76 percent for the Arkansas at Salida. Downstream, flow of the Arkansas at Pueblo should be about 67 percent. Flow of the Cucharas near LaVeta is also forecast at 67 percent.

With these low streamflow prospects and poor reservoir storage, some water shortages are expected during late summer months. The prospective shortages will become more severe if the present dry weather trend is not broken soon.

In New Mexico, the flow of the Canadian River should be a little more than one-half of average. Storage in Conchas Reservoir is 45 percent.

SELECTED STREAMFLOW FORECASTS

MAY 1, 1972

STREAM AND STATION	FORECASTS THIS YEAR		Forecast Period	Last Year's Flow In (1,000 A. F.)
	Flow in (1,000 A. F.)	Percent of Average		
SASKATCHEWAN				
St. Mary near Babb, Montana <u>1/</u>	565	120	May-Sept.	
UPPER MISSOURI				
Beaverhead near Grant, Montana <u>2/</u>	115	161	May-Sept.	248
Big Hole near Melrose, Montana	880	144	May-Sept.	
Jefferson at Sappington, Montana	1,230	151	May-Sept.	
Madison near Grayling, Montana <u>3/</u>	545	145	May-Sept.	619
Gallatin near Gateway, Montana	505	115	May-Sept.	700
Sun at Gibson Dam, Montana <u>4/</u>	780	136	May-Sept.	701
Belt near Monarch, Montana	147	143	May-Sept.	
Marias near Shelby, Montana <u>5/</u>	750	141	May-Sept.	520
Missouri near Landusky, Montana <u>6/</u>	5,600	142	May-Sept.	4,821
near Williston, North Dakota <u>7/</u>	13,500	140	May-Sept.	
S. Fk. Musselshell above Martinsdale, Montana	52.5	125	May-Sept.	45
Milk at Eastern Crossing, Montana	200	91	May-Sept.	190
Yellowstone at Yellowstone Lake Outlet, Wyo.	1,000	120	April-Oct.	1,217
at Corwin Springs, Montana	2,340	130	May-Sept.	2,607
at Miles City, Montana <u>8/</u>	7,400	136	May-Sept.	
Clarks Fork near Belfry, Montana	740	132	May-Sept.	812
Shoshone below Buffalo Bill Res., Wyo. <u>9/</u>	1,070	132	April-Sept.	1,150
Wind near Dubois, Wyoming	141	142	April-Sept.	
at Riverton, Wyoming <u>10/</u>	870	134	April-Sept.	
below Boysen Res., Wyoming <u>11/</u>	1,020	135	April-Sept.	
Bull Lake Creek near Lenore, Wyoming	236	133	April-Sept.	248
Little Popo Agie near Lander, Wyoming	59	139	April-Sept.	73
Tensleep near Tensleep, Wyoming	89	120	April-Sept.	88
Medicine Lodge near Hyattville, Wyoming	26	131	April-Sept.	21
Shell Creek near Shell, Wyoming	84	128	April-Sept.	
Big Horn near St. Xavier <u>8/</u>	2,300	144	May-Sept.	2,276
Tongue near Dayton, Wyoming	125	121	April-Sept.	112
No. Fork Powder near Hazelton, Wyoming	11	118	April-Sept.	11
PLATTE				
North Platte at Saratoga, Wyoming	560	101	April-Sept.	
Encampment near Encampment, Wyoming	140	110	April-Sept.	221
Laramie near Glendevy, Wyoming <u>12/</u>	55	90	April-Sept.	
Big Thompson at Drake, Colorado <u>13/</u>	90	90	April-Sept.	
Clear at Golden, Colorado <u>14/</u>	105	88	April-Sept.	
St. Vrain at Lyons, Colorado <u>15/</u>	60	86	April-Sept.	
Cache La Poudre near Fort Collins, Colorado <u>16/</u>	210	98	April-Sept.	
ARKANSAS				
Arkansas at Salida, Colorado <u>17/</u>	235	76	April-Sept.	
Cucharas near LaVeta, Colorado	8	67	April-Sept.	
Purgatoire at Trinidad, Colorado	28	60	April-Sept.	
RIO GRANDE				
Rio Grande near Del Norte, Colorado <u>18/</u>	300	68	April-Sept.	
at Otowi Bridge, New Mexico <u>19/</u>	260	51	March-July	
Conejos near Mogote, Colorado <u>20/</u>	110	60	April-Sept.	
El Vado Res., Inflow, New Mexico	120	64	March-July	
Pecos at Pecos, New Mexico	22	54	March-July	
UPPER COLORADO				
Colorado, Grandby Res. Inflow, Colorado <u>21/</u>	195	89	April-Sept.	
near Dotsero, Colorado <u>22/</u>	1,200	87	April-Sept.	
near Cameo, Colorado <u>23/</u>	1,850	83	April-Sept.	
near Cisco, Utah <u>24/</u>	1,884	67	April-July	
Lake Powell Inflow, Arizona <u>25/</u>	5,610	86	April-July	8,378
Roaring Fork at Glenwood Springs, Colorado <u>26/</u>	485	70	April-Sept.	
Uncompahgre at Colona, Colorado	80	62	April-Sept.	

Forecasts in California provided by Department of Water Resources.
Average is for 1953-67 period except California. California is computed for 1916-65 period.
Forecasts assume average Effective Climate Conditions from Date Through Snow Melt Season.

Explanatory Notes on Forecasts listed on Inside Back Cover.

SELECTED STREAMFLOW FORECASTS

MAY 1, 1972

STREAM AND STATION	FORECASTS THIS YEAR		Forecast Period	Last Year's Flow In (1,000 A.F.)
	Flow In (1,000 A.F.)	Percent of Average		
UPPER COLORADO (continued)				
Gunnison, Blue Mesa Res. Inflow, Colorado <u>27/</u>	500	65	April-Sept.	
near Grand Junction, Colorado <u>28/</u>	700	62	April-Sept.	
Dolores at Dolores, Colorado	150	65	April-Sept.	
Green at Warren Bridge, Wyoming	400	124	April-Sept.	452
at Green River, Wyoming <u>29/</u>	1,320	140	April-Sept.	
Flaming Gorge Res. Inflow, Utah <u>27/</u>	1,510	143	April-July	1,905
at Green River, Utah <u>30/</u>	2,964	115	April-July	
North Piney at Mason, Wyoming	55	160	April-Sept.	67
Big Sandy near Big Sandy, Wyoming	70	132	April-Sept.	69
Yampa at Steamboat Springs, Colorado	225	87	April-Sept.	
near Maybell, Colorado	750	88	April-Sept.	
Little Snake near Dixon, Wyoming	272	105	April-Sept.	486
White near Meeker, Colorado	210	72	April-Sept.	
Strawberry at Duchesne, Utah <u>40/</u>	40	100	May-July	
Duchesne near Tabiona, Utah <u>31/</u>	115	135	May-July	
at Randlett, Utah <u>40/</u>	260	114	May-July	
Lakefork below Moon Lake, Utah <u>32/</u>	60	95	May-July	
Uinta near Neola, Utah	73	97	May-July	
Whiterocks near Whiterocks, Utah	45	94	May-July	
Price, Scofield Res. Inflow, Utah <u>33/</u>	20	74	May-July	
Cottonwood near Orangeville, Utah <u>34/</u>	29	71	May-July	
San Juan, Navajo Res. Inflow, New Mexico <u>27/</u>	360	58	April-July	305
near Bluff, Utah <u>35/</u>	472	53	April-July	
Animas at Durango, Colorado	250	61	April-Sept.	
LOWER COLORADO				
Virgin near Virgin, Utah	10	45	May-June	
Little Colorado above Lyman, Arizona	---	---	---	---
Gila near Solomon, Arizona	---	---	---	---
Frisco at Clifton, Arizona	---	---	---	---
Salt Lake Intake, Arizona	25	21	April-May	27.1
Tonto above Roosevelt, Arizona	1.1	14	April-May	1.3
Verde above Horseshoe Dam, Arizona	18	36	April-May	21.0
GREAT BASIN				
Bear at Utah-Wyo. State Line	140	140	May-July	129
at Harer, Idaho	340	189	May-Sept.	
Smith's Fork near Border, Wyoming	157	145	April-Sept.	198
Thomas Fork near Wyo.Ida. State Line	51	162	April-Sept.	70
Logan near Logan, Utah <u>36/</u>	150	174	May-July	178
Ogden, Pine View Res. Inflow, Utah <u>27/</u>	102	173	May-June	92
Weber near Oakley, Utah	105	126	May-June	
Provo near Hailstone, Utah <u>37/</u>	98	120	May-July	
Strawberry Res. Inflow, Utah	26	83	May-July	
Utah Lake Net Inflow, Utah	160	118	May-July	
Big Cottonwood near Salt Lake City, Utah	36	120	May-July	
Beaver near Beaver, Utah	7.5	45	May-July	
Sevier near Hatch, Utah	15.3	57	May-July	
near Gunnison, Utah	10	46	May-July	
So. Fork Humboldt near Elko, Nevada	45	90	May-July	116
Humboldt at Palisades, Nevada	120	98	May-July	396
Truckee at Farad, California <u>38/</u>	150	79	May-July	309
East Carson near Gardnerville, Nevada	113	79	May-July	224
West Carson at Woodsfords, California	32	80	May-July	49
East Walker near Bridgeport, California <u>39/</u>	38	70	May-August	72
West Walker near Coleville, California	103	82	May-July	137
Donner und Blitzen near Frenchglen, Oregon	56	124	May-Sept.	
Silvies near Burns, Oregon	50	123	May-Sept.	58
Chewaucan near Paisley, Oregon	62	100	May-Sept.	129
Deep above Adel, Oregon	53	120	May-Sept.	102
Bidwell near Ft. Bidwell, California	15.3	170	May-July	14

Forecasts in California provided by Department of Water Resources.
Average is for 1953-67 period except California. California is computed for 1916-65 period.
Forecasts assume average Effective Climate Conditions from Date Through Snow Melt Season.

Explanatory Notes on Forecasts listed on Inside Back Cover.

SELECTED STREAMFLOW FORECASTS

MAY 1, 1972

STREAM AND STATION	FORECASTS THIS YEAR		Forecast Period	Last Year's Flow In (1,000 A.F.)
	Flow In (1,000 A.F.)	Percent of Average		
UPPER COLUMBIA				
Columbia above Steamboat Rapids, B. C.	21,500	122	May-Sept.	17,980
at Birchbank, British Columbia <u>40/</u>	53,570	123	May-Sept.	45,730
at Grand Coulee, Washington <u>40/</u>	80,000	127	May-Sept.	68,303
Kootenai at Libby, Montana	9,550	128	May-Sept.	8,354
at Leonia, Idaho	11,000	131	May-Sept.	9,662
Blackfoot near Bonner, Montana	1,330	148	May-Sept.	1,136
So. Fk. Flathead nr Columbia Falls, Montana <u>40/</u>	2,900	137	May-Sept.	2,521
Flathead at Columbia Falls, Montana <u>40/</u>	8,000	136	May-Sept.	6,878
near Polson, Montana <u>40/</u>	9,500	137	May-Sept.	7,989
Clark Fork above Missoula, Montana	2,370	153	May-Sept.	1,824
near Plains, Montana <u>40/</u>	16,000	144	May-Sept.	13,337
at Whitehorse Rapids, Idaho	17,800	144	May-Sept.	15,006
Bitterroot near Darby, Montana	755	150	May-Sept.	700
Priest near Priest River, Idaho <u>41/</u>	715	99	May-July	
Pend Oreille below Box Canyon, Washington	19,800	143	May-Sept.	
Kettle near Laurier, Washington	1,950	117	May-Sept.	2,020
Spokane at Post Falls, Idaho <u>42/</u>	3,500	166	May-Sept.	2,862
Similkameen near Nighthawk, Washington	2,080	147	May-Sept.	
Okanogan near Tonasket, Washington	2,450	152	May-Sept.	
Methow near Pateros, Washington	1,470	152	May-Sept.	
Stehekin at Stehekin, Washington	1,280	155	May-Sept.	
Chelan at Chelan, Washington <u>43/</u>	1,720	150	May-Sept.	
Wenatchee at Peshastin, Washington	2,409	150	May-Sept.	
SNAKE				
Snake above Palisades Res., Wyoming <u>44/</u>	3,520	138	April-Sept.	4,048
near Heise, Idaho <u>45/</u>	4,650	136	May-Sept.	5,766
near Blackfoot, Idaho <u>46/</u>	4,750	135	May-July	
at Weiser, Idaho	6,250	125	May-Sept.	8,300
Grey's above Palisade, Wyoming	590	163	April-Sept.	
Salt above Palisade, Wyoming	500	156	April-Sept.	
Henry's Fork near Ashton, Idaho <u>47/</u>	565	110	May-Sept.	
Teton near St. Anthony, Idaho	405	115	May-Sept.	
Blackfoot Reservoir Inflow, Idaho	135	132	April-Sept.	
Big Lost near MacKay, Idaho <u>48/</u>	165	104	May-Sept.	
Portneuf at Topaz, Idaho	70	125	May-Sept.	
Salmon Falls Creek nr San Jacinto, Idaho	70	151	May-Sept.	
Big Wood, Inflow to Magic Res., Idaho <u>49/</u>	260	141	May-Sept.	
Bruneau near Hot Springs, Idaho	210	148	May-Sept.	
Boise near Boise, Idaho <u>50/</u>	1,800	147	May-Sept.	2,087
Jordan near Jordan Valley, Oregon	66	138	May-July	
Owyhee near Owyhee, Nevada <u>51/</u>	50	132	May-July	128
Owyhee Res. Net Inflow, Oregon <u>27/</u>	232	130	May-Sept.	277
Malheur near Drewsey, Oregon	53	156	May-Sept.	61
Payette near Horseshoe Bend, Idaho <u>52/</u>	1,950	129	May-Sept.	2,485
Weiser above Crane Creek, Idaho <u>40/</u>	325	122	May-Sept.	
Burnt near Hereford, Oregon <u>40/</u>	20.4	132	May-Sept.	29
Powder near Sumpter, Oregon	44	118	May-Sept.	
Eagle above Skull Creek, Oregon	210	135	May-Sept.	240
Imnaha at Imnaha, Oregon	300	133	May-Sept.	380
Salmon at Whitebird, Idaho	8,050	130	May-Sept.	9,682
Lostine near Lostine, Oregon	148	128	May-Sept.	151
Grande Ronde at LaGrande, Oregon	120	114	May-Sept.	126
Clearwater at Spalding, Idaho	10,250	150	May-Sept.	9,067
LOWER COLUMBIA				
Yakima at CleElum, Washington <u>53/</u>	1,217	154	May-Sept.	
near Parker, Washington <u>54/</u>	2,158	165	May-Sept.	
Naches near Naches, Washington <u>55/</u>	1,197	160	May-Sept.	
Walla Walla, So. Fk. near Milton, Oregon	65	130	May-Sept.	68

Forecasts in California provided by Department of Water Resources.

Average is for 1953-67 period except California. California is computed for 1916-65 period.

Forecasts assume average Effective Climate Conditions from Date Through Snow Melt Season.

Explanatory Notes on Forecasts listed on Inside Back Cover.

SELECTED STREAMFLOW FORECASTS


MAY 1, 1972

STREAM AND STATION	FORECASTS THIS YEAR		Forecast Period	Last Year's Flow In (1,000 A.F.)
	Flow In (1,000 A.F.)	Percent of Average		
LOWER COLUMBIA (continued)				
Umatilla at Pendleton, Oregon	91	113	May-Sept.	88
John Day, Middle Fork at Ritter, Oregon	80	108	May-Sept.	88
North Fork at Monument, Oregon	376	100	May-Sept.	
Crooked near Post, Oregon	52	130	May-Sept.	
Deschutes at Benham Falls, Oregon <u>40/</u>	609	119	May-Sept.	
Columbia at The Dalles, Oregon <u>40/</u>	118,500	128	May-Sept.	108,712
Hood near Tucker Bridge, Oregon <u>40/</u>	295	121	May-Sept.	
McKenzie near Vida, Oregon	1,312	133	May-Sept.	
Santiam, South, at Waterloo, Oregon	506	135	May-Sept.	
North, at Mehama, Oregon <u>40/</u>	754	123	May-Sept.	
Clackamas at Estacada, Oregon	709	125	May-Sept.	
Willamette at Salem, Oregon <u>40/</u>	4,075	124	May-Sept.	
Lewis at Ariel, Washington <u>56/</u>	1,410	147	May-Sept.	1,397
Cowlitz at Castle Rock, Washington <u>57/</u>	2,940	139	May-Sept.	
NORTH PACIFIC COASTAL				
Dungeness near Sequim, Washington	166	108	May-Sept.	
Umpqua, No., near Tokatee Falls, Oregon <u>40/</u>	165	112	May-Sept.	
Rogue at Raygold, Oregon	740	108	May-Sept.	961
Klamath Lake, Net Inflow, Oregon	419	100	May-Sept.	563
Trinity at Lewiston, California	460	75	April-July	734
CALIFORNIA CENTRAL VALLEY <u>40/</u>				
Sacramento, Inflow to Shasta, California	1,530	86	April-July	2,332
Feather near Oroville, California	1,180	63	April-July	2,701
Yuba at Smartville, California	750	70	April-July	1,387
American, Inflow to Folsom Res., Calif.	840	64	April-July	1,445
Cosumnes at Michigan Bar, California	60	41	April-July	123
Mokelumne, Inflow to Pardee Res., Calif.	300	65	April-July	490
Stanislaus, Inflow to Melones Res., Calif.	420	59	April-July	664
Tuolumne, Inflow to Don Pedro Res., Calif.	730	61	April-July	1,058
Merced, Inflow to Exchequer Res., Calif.	310	51	April-July	502
San Joaquin, Inflow to Millerton Lake, Calif.	580	49	April-July	970
Kings, Inflow to Pine Flat Res., California	445	38	April-July	820
Kaweah, Inflow to Terminus Res., California	65	24	April-July	196
Tule, Inflow to Success Res., California	7	12	April-July	36.7
Kern, Inflow to Isabella Res., California	85	20	April-July	230
ALASKA				
Chena at Fairbanks, Alaska	750	170	May-June	658
Salcha near Salchaket, Alaska	855	146	May-June	878

Forecasts in California provided by Department of Water Resources.
Average is for 1953-67 period except California. California is computed for 1916-65 period.
Forecasts assume average Effective Climate Conditions from Date Through Snow Melt Season.

Explanatory Notes on Forecasts listed on Inside Back Cover.

RESERVOIR STORAGE as of MAY 1, 1972



RIO GRANDE BASIN

Only the extreme high elevations in New Mexico have any snow left. On the Rio Grande headwaters in Colorado the snow is only a little above half of the usual amount. This poor snow cover indicates that flow of the Rio Grande River and its tributaries will be near 50 to 65 percent in New Mexico, 60 to 70 percent in Colorado. Flow of the Pecos River is expected to be essentially half of average (54 percent).

Unless summer rains are unusually plentiful, water supplies will be very deficient. This applies particularly to water users on natural flow rights and to those having limited storage rights. Storage in Elephant Butte Reservoir is 53 percent of average, while El Vado holds 26 percent of its normal supply. Carry-over storage for next year will be very poor.

Extreme conservation of water supplies will be necessary this summer.

COLORADO BASIN

Dry weather during April has reduced water supply prospects for the Colorado River Basin during the coming summer.

The present snow cover in the upper Colorado River Basin is near 85 percent of average when considered as a whole, but it shows extreme variability within the Basin. Snow cover is heaviest on tributaries to the Green River in

Wyoming where it averages 152 percent of usual amounts. The snow decreases steadily south of here, with about 75 to 95 percent shown on the Yampa, White, upper Colorado (in Colorado) and Duchesne rivers. It drops to 50 to 65 percent on the Gunnison, San Juan, Price and San Rafael rivers. It is only 15 percent on the Dolores River.

Forecasts for streams tributary to the upper Green River in Wyoming range from about 125 to 160 percent of average. Total inflow to Flaming Gorge Reservoir in Utah is now expected to be 143 percent. In Colorado the Yampa and upper Colorado should yield near 12 percent less than usual, while the White, Gunnison, Roaring Fork and Uncompahgre rivers should flow at near 60 to 70 percent, with the Animas and San Juan yielding about 50 to 60 percent. In Utah the Duchesne will supply about 15 percent more than usual, while the Price and San Rafael will fall 25 to 30 percent below normal.

Unregulated flow of the principal tributaries is forecast as follows: Green at Green River, Utah 115 percent; Colorado near Cisco, Utah 67 percent and San Juan near Bluff, Utah 53 percent. Total inflow to Lake Powell, Arizona is forecast at 5,610,000 acre-feet for the April-July period, or 86 percent average. Storage in irrigation reservoirs is well above average.

April weather continued its dry pattern in the Lower Colorado Basin. All watersheds have prospects for below normal streamflow. Highest streamflow forecast is for the Virgin River in Utah (45 percent). Snowmelt runoff in

STORAGE IN LARGE RESERVOIRS

MAY 1, 1972

BASIN AND NAME OF RESERVOIR	CAPACITY (1,000 A.F.)	STORAGE (1,000 A.F.)	STORAGE PERCENT AVERAGE	BASIN AND NAME OF RESERVOIR	CAPACITY (1,000 A.F.)	STORAGE (1,000 A.F.)	STORAGE PERCENT AVERAGE
UPPER MISSOURI				UPPER COLUMBIA			
Belle Fourche	185	165	148	Chelan	676	205	98
Boysen	550	252	71	Coeur d'Alene	225	209	73
Buffalo Bill	373	106	86	Duncan	1,347	52	---
Canyon Ferry	2,043	1,533	98	Flathead	1,791	976	104
Fort Peck	19,410	16,550	148	Hungry Horse	3,428	1,255	64
Garrison	24,790	21,380	192	Kootenay	673	189	41
Hebgen	377	182	93	Lower Arrow	3,083	31	6
Keyhole	192	169	433	Noxon Rapids	335	182	126
Lake Francis Case	5,816	4,487	115	Pend Oreille	1,155	346	70
Lake Sharp	1,900	1,743	104	Roosevelt	5,232	497	20
Oahe	23,630	19,899	156	Upper Arrow	4,061	6	1
Tiber	1,347	558	85	LOWER COLUMBIA			
Big Horn	1,356	792	107	Cougar	155	129	---
PLATTE				Detroit	300	255	110
City of Denver (5)	507	425	111	Green Peter	270	208	---
Colo-Big Thompson (3)	718	560	137	Hills Creek	200	165	101
Glendo	784	502	124	Lookout Point	337	246	85
Pathfinder	1,016	925	211	Prineville	153	152	103
Seminole	1,010	583	195	Wickiup	200	198	102
ARKANSAS				Yakima Res. (5)	1,066	511	62
Conchas	273	68	45	SNAKE			
John Martin	354	0	0	American Falls	1,700	1,650	99
RIO GRANDE				Anderson Ranch	423	254	89
Elephant Butte	2,195	172	53	Arrowrock	287	142	62
El Vado	195	8	26	Brownlee	980	21	---
UPPER COLORADO				Cascade	653	218	67
Blue Mesa	830	320	---	Jackson	847	501	114
Flaming Gorge	3,749	2,728	---	Lucky Peak	278	56	38
Navajo	1,696	847	---	Owyhee	715	708	133
Powell	25,002	13,224	---	Palisades	1,200	199	25
Starvation	152	127	---	Warm Springs	191	180	131
LOWER COLORADO				PACIFIC COASTAL			
Havasu	619	592	100	Clair Engle	2,448	2,349	107
Mead	26,159	17,015	106	Clear Lake	440	416	156
Mohave	1,810	1,689	98	Macimiento	350	62	28
Salt River Res. (4)	1,755	870	84	Ross	1,203	769	110
San Carlos	985	82	73	Upper Klamath	584	538	104
Verde River Res. (2)	318	64	47	CALIFORNIA CENTRAL VALLEY			
GREAT BASIN				Almanor	1,036	782	98
Bear	1,421	1,204	126	Berryessa	1,602	1,361	87
Lahontan	314	277	125	Folsom	1,010	865	120
Rye Patch	179	188	227	Isabella	570	113	58
Sevier Bridge	236	173	183	McClure	1,026	574	91
Strawberry	274	219	166	Millerton	521	338	93
Tahoe	732	582	126	Bullards Bar	930	757	117
Utah	884	856	138	Oroville	3,484	3,424	124
Willard Bay	193	177	---	Pine Flat	1,013	489	73
				Shasta	4,500	4,363	103

Reservoir Storage Data Provided by Bureau of Reclamation, Corps of Engineers, Geological Survey, and water using organizations. Data from California and British Columbia provided by Department of Water Resources and Department of Lands, Forests and Water Resources, respectively.

Arizona has occurred. The expected runoff of 15 to 35 percent is base flow on most streams.

Fortunately, reservoir storage, though steadily declining, is 80 percent of average for this date. There is adequate water for this year on most projects, but carryover will be very low and severe shortages are possible next year. Heavy ground water pumping will again be required in the Gila Valley to meet water demands.

GREAT BASIN

Considering the very favorable reservoir storage throughout the Great Basin, with streamflow prospects for most major irrigated areas varying from near 75 percent to more than double usual amounts, the summer's water supply should prove to be fair to exceptionally good for most water users.

Late summer shortages are anticipated in Nevada on smaller streams coming from lower elevation watersheds, particularly in southern areas. In southern Utah on the Sevier, San Pitch and Beaver rivers and smaller streams near Cedar City, water users on natural flow rights will experience definite shortages. Flow of these streams is expected to range from about 45 to 70 percent of average. Fortunately, storage in the three main Sevier River reservoirs is 168 percent of average.

The snowpack varies considerably. In southern areas of Utah and Nevada the snow ranges from zero to near half of average. In northern Utah and on Nevada's Jarbidge Mountains snow water equivalents vary from average to 175 percent. In Nevada's Sierra Nevada Mountains the low and mid-elevation snowpack was depleted early. The remaining high elevation snowpack is normal for this date.

Highest May-July runoff forecasts in Nevada are on northern tributaries to the Humboldt River near Elko. These streams are expected to produce 20 to 40 percent above normal. The main Humboldt and its other tributaries are forecast at 90 to 110 percent, flow of the Truckee, Carson, Walker and Owens rivers will be near 70 to 80 percent. Reservoirs in Nevada now hold 86 percent of their capacity and are at 140 percent of normal.

Flow of the Bear River and its tributary streams in Utah, Idaho and Wyoming will generally be near 140 percent to over twice their usual amounts. Streamflow on the Weber-Ogden rivers will be similar to the Bear River. Prospective flow of streams draining into Utah Lake varies from about 80 to 120 percent.

Reservoir storage in 14 of Utah's major

reservoirs was 138 percent of the May 1 average for the 1953-67 period.

In southern Oregon's Harney County the water outlook is very favorable, with streams forecast at 20 to 25 percent above normal. In Lake County the forecasts range from average to 20 percent above.

In view of the water outlook in surrounding areas, prospects for California's Surprise Valley are surprising. Streams here are forecast at 170 to 185 percent of normal amounts.

COLUMBIA BASIN

The water supply outlook remains good to excellent throughout the Columbia Basin. Cool April weather delayed snowmelt and maintained a serious flood potential on many watersheds, particularly if an adverse sequence of temperatures and/or precipitation should develop during the main snowmelt period.

Most watersheds in the Columbia Basin have a snowpack which is over 140 percent of average. Percentagewise, the snow is particularly heavy along the Cascade Mountains of Washington, British Columbia and the Hood River of Oregon; on the east Kootenay in British Columbia, Montana's upper Clark Fork and Bitterroot rivers, Idaho's Clearwater and Bruneau rivers and Oregon's Owyhee, Malheur and Umatilla rivers. The snow on these streams ranges from near 165 to over 200 percent of average.

In the above areas, many snow courses have maximum or near maximum of record snowpacks for May 1st. Some courses have set new all season record high readings.

The present heavy snowpack indicates that the main snowmelt period will produce an unregulated peak flow of near 850 thousand to one million second feet at The Dalles, Oregon. These flows would result in a river stage at Vancouver of 28 to 31 feet. Actual flow and river stage will depend on snowmelt conditions during late May and early June, and on the amount that river regulation reduces the flow. River management agencies have indicated that reservoir regulation will reduce the river stage at Vancouver to about 21 to 23 feet.

Most major irrigation reservoirs are full or nearly full, except where they have been drawn down to provide space for flood control. These should refill and reservoir storage carryover for next year should be excellent.

Because of the heavy snowpacks, record or near record high streamflow is expected from streams in British Columbia, Montana, the

Spokane and Clearwater rivers in Idaho and streams flowing from the Cascades in Washington. Forecasts for most of these streams range from about 130 to 165 percent of normal.

Although not record or near record high, forecasts for most tributaries to the Snake River are far above average and vary from about 125 to 160 percent. Lowest forecast on the Snake River is 95 percent for Idaho's Little Wood River, while the Big Wood, Henry's Fork, Teton, Grande Ronde, Weiser and Powder rivers are forecast at 5 to 20 percent above normal. Most Oregon streams other than those that are tributary to the Snake River will yield from near average to 30 percent above average amounts.

ALASKA

Many snow courses in Alaska have a snowpack which is maximum of record. Cool April weather left the low elevation snow unmelted, while the higher elevation snow continued to increase. Spring runoff will approach 150 to 200 percent of normal over a wide area of Alaska. Among these will be interior rivers such as the Yukon and Tanana, as well as southeastern Alaska streams, and rivers discharging into Cook Inlet.

CALIFORNIA

The California Department of Water Resources, coordinating agency for snow surveys in California, reports that the May 1 snow surveys along with reports on precipitation, streamflow, etc., confirm that only in watersheds along the Oregon border are normal conditions prevailing. Forecasts of snowmelt runoff for the Central Valley tributaries indicate 1972 April-July flows will be the sixth lowest of the past 20 years. Although this summer will be dry and probably hot, no general critical water shortages are expected due to ample supply of storage in surface reservoirs along with facilities to transfer large amounts to areas of need plus the ability to supplement surface supplies by pumping from ground water sources.

In Southern California, the benefit to local supplies received from the heavy precipitation of last December has fast eroded away by subsequent months of record drought. Thus, the dependence upon local supplies becomes a way of life.

Precipitation in California during April mostly came in the first half of the month from two moderate to light storms. Statewide, precipitation for April averaged 80 percent of normal with above normal amounts generally

limited to the central Sierra and North Coastal area. In Southern California, April produced some 0.03 inch of precipitation at the Los Angeles Civic Center which made a total for the 4-month period January through April of 0.16 inch at this station, an all-time low and some 9.50 inches below that normally expected. May 1 generally marks the end of the major precipitation and snow accumulation in California. While it is still possible for rather heavy general storms to materialize, such occurrences cannot be planned upon.

May 1 snow surveys of some 160 key snow courses and 65 aerial marker observations indicate that the water content in the State's snowpack was about 70 percent of normal for May 1 or 50 percent of the April 1 average. Although many high elevation snow courses from the Kings River Basin north showed water content gains since April 1, most low and many middle elevation snow courses were bare. The May 1 measurements indicated that only in the Central Sierra was the snowpack's water content near normal. To the north, the Feather River Basin dropped to 70 percent of normal while to the south the Tule River Basin was only 5 percent of the normal for May 1.

Runoff from California's major streams during April was below average over the entire State, averaging about 75 percent of normal. Runoff from nonsnowmelt streams during April averaged less than 15 percent of normal, except in the North Coastal area where runoff for the month was 88 percent of normal. Runoff from snowmelt streams tributary to the Central Valley averaged about 70 percent of normal for the month. In the Sacramento Valley, the April runoff for individual river basins ranged from slightly below normal for the Upper Sacramento River to 54 percent of normal for the Cosumnes River Basin. San Joaquin Valley tributaries' runoff varied from a high of 63 percent of normal for the Stanislaus River Basin to a low of 16 percent of normal for the Tule River Basin.

Storage on May 1, 1972 in 121 major California reservoirs, with a combined capacity of 29,824,000 acre-feet, was 21,332,000 acre-feet, slightly above the May 1 average for the aggregate storage. This is just over 1,000,000 acre-feet less than was in storage at this time one year ago. In the Central Valley, aggregate storage in the major reservoirs was just average for May 1. With variation within the individual river basins, aggregate storage in Sacramento and San Joaquin Valley major reservoirs was 105 and 95 percent of their 10-year average, respectively.



EXPLANATION of STREAMFLOW FORECASTS

All flows are observed flows except as adjusted for: 1/ Storage change in Lake Shoshone. 2/ Storage change in Lima and Clark Canyon reservoirs. 3/ Storage change in Hebgen Lake. 4/ Storage change in Gibson Reservoir and measured diversions. 5/ Storage change in Two Medicine, Four Horns, Lake Francis and Swift reservoirs. 6/ Storage change in Canyon Ferry and Tiber reservoirs. 7/ Changes as indicated in (6/), (8/), plus storage change in Fort Peck. 8/ Storage change in Boysen, Buffalo Bill and Yellowtail reservoirs. 9/ Storage change in Buffalo Bill Reservoir plus Heart Mountain diversion. 10/ Storage change in Pilot Butte and Bull Lake reservoirs plus Wyoming canal diversion.

11/ Changes indicated in (10/) plus storage change in Boysen Reservoir. 12/ Plus diversions to Cache LaPoudre. 13/ Plus by-pass to power plants. 14/ Minus diversion thru Gumlick Tunnel. 15/ Storage change in Price Reservoir. 16/ Minus diversions from North Platte, Laramie and Colorado rivers plus measured diversions above station. 17/ Storage change in Clear Creek, Twin Lakes and Turquoise reservoirs minus diversions from Colorado River. 18/ Storage change in Rio Grande, Santa Maria and Continental reservoirs. 19/ Storage change in El Vado and Abiquiu reservoirs. 20/ Storage change in Platoro Reservoir.

21/ Storage change in Grandby Reservoir as furnished by U.S.B.R. plus diversions by Adams Tunnel and Grand River Ditch. 22/ Changes as indicated in (21/) plus diversions thru Roberts, Gumlick and Moffat tunnels and storage change in Dillon, Homestake, Williams Fork, Green Mountain and Willow Creek reservoirs. 23/ Changes indicated in (22/) and (26/). 24/ Storage change in Blue Mesa Reservoir. 25/ Changes indicated in (24/), (30/) and (35/) and storage change in Lake Powell. 26/ Diversions to Arkansas River plus storage change in Ruedi Reservoir. 27/ (Inflow record as computed by U. S. Bureau of Reclamation.) 28/ Storage change in Taylor, Blue Mesa and Morrow Point reservoirs. 29/ Storage change in Fontenelle Reservoir. 30/ Storage change in Flaming Gorge Reservoir.

31/ Plus diversion through Duchesne Tunnel. 32/ Storage change in Moon Lake Reservoir. 33/ Storage change in Scofield Reservoir. 34/ Storage change in Joe's Valley Reservoir. 35/ Storage change in Navajo Reservoir. 36/ Plus U. P. & L. Co. tailrace and Logan, Hyde Park and Smithfield canals. 37/ Minus diversions thru Duchesne Tunnel and Weber-Provo Canal. 38/ Storage change in Lake Tahoe and Boca reservoirs (Forecast by Truckee Basin Committee.) 39/ Storage change in Bridgeport Reservoir. 40/ Corrected for major upstream impairments -- represents simulated natural flow conditions.

41/ Storage change in Priest Lake. 42/ Storage change in Coeur d'Alene Lake and diversions by Spokane Valley Farms Co. and Rathrum Prairie canals. 43/ Storage change in Lake Chelan. 44/ Storage change in Jackson Lake. 45/ Storage change in Jackson Lake and Palisade reservoirs. 46/ Storage change in Jackson Lake, Palisades, Island Park, Henry's Lake, Grassy Lake plus diversions between Heise and Blackfoot. 47/ Storage change in Henry's Lake and Island Park reservoirs. 48/ Storage change in MacKay Reservoir and diversion in Sharp Ditch. 49/ Combined flow Big Wood near Bellevue and Camas Creek near Blaine. 50/ Storage change in Arrowrock, Anderson Ranch and Lucky Peak reservoirs.

51/ Storage change in Wild Horse Reservoir. 52/ Storage change in Cascade and Deadwood reservoirs. 53/ Storage change in Keechelus, Kachess and CleElum reservoirs plus diversion by Kittitas Canal. 54/ Changes indicated in (52/) plus storage change in Bumping and Rimrock Lakes plus diversion by Roza, Union Gap, New Reservation, Old Reservation and Sunrise canals. 55/ Storage change in Bumping and Rimrock lakes and diversions by Tieton, Selah Valley, Wapatox canals and City of Yakima. 56/ Storage change in Merwin, Yale and Swift reservoirs. 57/ Storage change in Mayfield Reservoir.

UNITED STATES DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE
701 N.W. GLISAN, RM. 209
PORTLAND, OREGON 97209

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE, \$300

POSTAGE AND FEES PAID
U. S. DEPARTMENT OF
AGRICULTURE



FIRST CLASS MAIL

FEDERAL - STATE - PRIVATE
COOPERATIVE SNOW SURVEYS

Furnishes the basic data
necessary for forecasting
water supply for irrigation,
domestic and municipal water
supply, hydro-electric power
generation, navigation,
mining and industry

*"The Conservation of Water begins
with the Snow Survey"*